

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> :**A61B 19/00****A1**

(11) International Publication Number:

**WO 98/38942**

(43) International Publication Date: 11 September 1998 (11.09.98)

(21) International Application Number: PCT/US98/04797

(22) International Filing Date: 18 February 1998 (18.02.98)

(30) Priority Data:

08/812,879

6 March 1997 (06.03.97)

US

(71) Applicant: SCIMED LIFE SYSTEMS, INC. [US/US]; One  
SciMed Place, Maple Grove, MN 55311-1566 (US).(72) Inventors: RESSEMANN, Thomas, V.; 1730 Boxwood Circle,  
St. Cloud, MN 56303 (US). HARRISON, Kent, D.; 15407  
Elm Road North, Maple Grove, MN 55311 (US).(74) Agents: KELLY, Joseph, R.; Westman, Champlin & Kelly,  
P.A., International Centre - Suite 1600, 900 Second Avenue  
South, Minneapolis, MN 55402-3319 (US) et al.(81) Designated States: CA, JP, European patent (AT, BE, CH, DE,  
DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).


Published

*With international search report.*

(54) Title: PERCUTANEOUS BYPASS BY TUNNELING THROUGH VESSEL WALL

## (57) Abstract

A parent vessel (16) lies adjacent heart tissue, and a restriction (24) in the parent vessel (16) is bypassed. A first region of the parent vessel (16) on a first side of the restriction (24) is accessed and an aperture is formed therein. A lumen (40) is formed through the heart tissue, and the lumen (40) communicates with the parent vessel (16) through the first aperture. A second aperture is formed in a second region of the parent vessel (16) on a second side of the restriction (24). The lumen (40) through the heart tissue communicates with the parent vessel (16) through the first and second apertures, thus bypassing the restriction (24).



*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-2-

for use as the bypass graft. One common source for the bypass graft is the saphenous vein which is removed from the leg. Harvesting the saphenous vein requires the surgeon to cut and peel the skin back from an area of the leg which is approximately 18 inches long and which extends upward to the groin area. This can be very traumatic and painful. Also, the internal mammary artery (IMA) has also been used as a vein graft in performing a bypass. However, the IMA is typically best suited for use as a left anterior descending (LAD) vein graft and is commonly saved for that purpose. Further, open heart surgery requires quite a lengthy recovery period which involves an increased hospital stay, and, consequently, greater expense.

Other than the pain and more lengthy hospital stay, open heart surgery involves other disadvantages as well. For example, during open heart surgery, it is common to cool the heart to a point where it stops. The blood from the remainder of the vasculature is then pumped through a pulmonary and cardiac bypass system. Any time the heart is stopped, there is a danger of encountering difficulty in restarting the heart (which is typically accomplished by warming the heart and massaging it). Further, even if the heart is restarted, it sometimes does not return to a correct rhythm. Also, open heart surgery can require the use of a device known as a left ventricular assist device (LVAD) to supplementarily pump blood to relieve the burden on the heart. This allows the heart to heal.

A significant reason that the heart is typically stopped during open heart surgery is that, if it were not stopped, the surgeon would be working in a dynamic environment. In such an environment, the target vessels and tissue to be treated are moving. Further,

-3-

a system must be employed in such an environment to stop bleeding. Clinical studies indicate that, when blood flow is stopped using clamping devices and blood flow is diverted to a cardiac bypass system, a statistically significant instance of neurological problems caused by blood clotting results. The use of mechanical clamps to stop blood flow, and the use of a mechanical bypass system, results in an approximate six percent instance of neurological problems, such as stroke, memory failure, etc.

Given the difficulties of the techniques discussed above, another approach has been developed which does not require stoppage of the heart or an open chest during execution. This approach is to perform a bypass using a minimally invasive technique by entering the upper chest cavity, through a hole between ribs under visual observation. Such a technique is often referred to as minimally invasive direct coronary artery bypass (MIDCAB) (where the heart is not stopped) or heart port (where the heart is stopped). Such a system which is used to perform a bypass is disclosed in the Sterman et al. U.S. Patent No. 5,452,733.

#### SUMMARY OF THE INVENTION

A parent vessel lies adjacent heart tissue, and a restriction in the parent vessel is bypassed. A first region of the parent vessel on a first side of the restriction is accessed and an aperture is formed therein. A lumen is formed through the heart tissue, and the lumen communicates with the parent vessel through the first aperture. A second aperture is formed in a second region of the parent vessel on a second side of the restriction. The lumen through the heart tissue communicates with the parent vessel through the first and second apertures, thus bypassing the restriction.

-4-

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1A illustrates a portion of a coronary vascular system with an artery having a restriction.

5 FIG. 1B illustrates a greatly enlarged cross-sectional view of a portion of the heart and artery shown in FIG. 1A in the area of the restriction.

FIG. 2. illustrates a boring device accessing a region proximal of the restriction.

10 FIG. 3 illustrates the boring device forming a lumen through heart tissue

FIG. 4 illustrates the boring device shown in FIG. 3 extending the lumen through the heart tissue into a region of the artery distal of the restriction.

15 FIG. 5 illustrates deployment of a drug delivery device in the lumen through the heart tissue.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

FIG. 1A illustrates a portion of a vascular system 10 including heart 12, aorta 14, and an artery 16. FIG. 1 also illustrates a plurality chambers 18 and 20 in heart 12, and a valve 22. A restriction 24 is illustrated in artery 16.

FIG. 1B is a greatly enlarged cross-sectional view of a portion of heart 12, and vessel 16 in the area of restriction 24. FIG. 1B illustrates that restriction 25 24 is typically covered within vessel 16 by an endothelial cell layer 13. Also, the wall of artery 16, and the heart tissue adjacent the wall of artery 16 is made up of a plurality of layers including internal elastic lamina (IEL) 15, media 17, adventitia 19, and 30 myocardium 21. Typically, the depth of the IEL layer 15, media layer 17 and adventitia layer 19, together, is approximately 1 mm. Also, typically, the thickness of the myocardium layer 21 is approximately 5-10 mm. The present invention bypasses restriction 24 by tunneling

-5-

around restriction 24 through the layers of artery 16, and heart 12. It should specifically be noted that the present invention, when referring to boring or tunneling through heart tissue, contemplates that restriction 24  
5 can be bypassed by tunneling around it through myocardium layer 21, or short of myocardium layer 21, and through one or more of IEL layer 15, media layer 17 and adventitia layer 19. Thus, the path of the tunnel around restriction 24 need not pierce all the way to  
10 myocardium layer 21.

FIG. 2 illustrates a first step utilized in accordance with the present invention to bypass restriction 24 in artery 16. A standard guide catheter 26 is advanced through aorta 14 in a known manner.  
15 Guide catheter 26 is positioned such that its distal end 28 is located in a region proximate the ostium 30 of artery 16. Cutting device 32 is then advanced through guide catheter 26, out through distal end 28 of guide catheter 26, through ostium 30 of vessel 16 and to a  
20 region proximal of restriction 24 in vessel 16. Cutting device 32, in the preferred embodiment, includes a catheter portion 34 and a boring or cutting member 36. Boring or cutting member 36 can take any number of suitable forms, such as a rotatable member or blade  
25 which can be used in piercing and boring through tissue. It should be noted that cutting device 32 can be any suitable mechanical cutting device, or any suitable sort of energy device, such as a radio frequency (RF) ablation wire, or a laser, or other device.

30 FIG. 3 illustrates an additional step utilized in accordance with the present invention in bypassing restriction 24 in vessel 16. Cutting device 32 is advanced to the region proximal of restriction 24 in vessel 16. Cutting member 36 is then used to pierce the

-6-

wall of vessel 16 to form an aperture 38 in the region proximal of restriction 24. Alternatively, cutting device 32 can be used to create another ostium in aorta 14 preferably in a region proximate the ostium of vessel 16. In that case, cutting device 32 need not be advanced into vessel 16. In either case, cutting device 32 is then further advanced through the myocardium 21 (or only through any or all of the other tissue layers 15, 17 and 19) closely proximate vessel 16. In one preferred embodiment, cutting device 32 is advanced such that lumen 40 is generally parallel to lumen 16 in the region proximate restriction 24.

FIG. 4 illustrates yet another step utilized in accordance with one preferred embodiment of the present invention for bypassing restriction 24 in vessel 16. FIG. 4 illustrates that, in the preferred embodiment, cutting device 32 is advanced such that cutting tip 36 advances through lumen 40 in the heart tissue to again pierce the vessel wall of vessel 16 to form a second aperture 42 therein. Second aperture 42 is located distal of restriction 24. In this way, lumen 40 extends from aperture 38 (proximal of restriction 24) to aperture 42 (distal of restriction 24) such that lumen 40 bypasses restriction 24 in vessel 16.

In a preferred embodiment, aspiration is provided during the cutting operation to remove severed tissue pieces. Such aspiration is preferably accompanied by saline infusate to flush the area around cutting tip 36. Aspiration can be provided using any suitable known technique, such as an aspiration catheter.

FIG. 5 illustrates another step in accordance with one preferred embodiment of the present invention. In FIG. 5, cutting device 32 has been withdrawn from

-7-

vascular system 10 through guide catheter 26. Drug delivery device 44 has been advanced through guide catheter 28, through ostium 30 in vessel 16, through aperture 38 in vessel 16 and into lumen 40 through the heart tissue. Drug delivery device 48 is preferably any suitable and known drug delivery device, such as a microcatheter, a perforated balloon delivery device, or any other suitable device. In the preferred embodiment, drug delivery device 48 is used to deliver a drug to lumen 40 to enhance endothelial development in lumen 40. In one preferred embodiment, the substance delivered to lumen 40 with delivery device 48 includes a growth factor which enhances the growth of endothelial cells on the wall of lumen 40. In another preferred embodiment, the substance delivered to lumen 40 with delivery device 48 includes endothelial cells which enhance endothelialization in lumen 40.

Cutting device 32 can be guided through the heart tissue in any number of suitable ways from the point proximal of restriction 24 to the point distal of restriction 24 where it re-enters vessel 16. For instance, the cutting member 36 of cutting device 32 is preferably formed of radiopaque material or has radiopaque markings thereon. Therefore, when restriction 24 is not a total occlusion, contrast fluid is injected through vessel 16 to locate restriction 24. The relative position of the restriction 24 and cutting device 36 is then observed.

In another embodiment, radiopaque markers are placed in vessel 16 distal of restriction 24. Bi-plane fluoroscopy is then used to perform three dimensional visualization of the markers in vessel 16 and on cutting member 32 to bring them toward one another.



-8-

In another embodiment, restriction 24 is crossed with a conductive wire or fiber. Either the conductive wire or fiber residing in vessel 16 distal of restriction 24, or cutting device 32, are provided with a transmitter, and the other (either the wire or fiber which has been used to cross restriction 24, or cutting device 32) is provided with a receiver or sensor. In one preferred embodiment, the transmitter includes an array of active transmitters comprising one of ultrasound transmitters, radiofrequency transmitters, a plurality of point light sources, or a single intense point light source, or an electro-magnetic transmitter (such as where current is actively applied to a coil to induce a magnetic field thereabout). The receiver, or sensor, is a suitable device which is compatible with the transmitter so that it can receive or sense the signals provided by the transmitter.

For instance, where the transmitter includes an inductive magnetic coil, the receiver includes a magnetic sensor array to receive the signals induced in the coil. When the transmitter includes an ultrasound transmitter, the receiver includes an ultrasound imager so that the relative positioning of the receiver device in the transmitter can be determined. When the transmitter includes a single point light source, or an array of point light sources, the receiver or sensor includes a photodiode array or an imaging fiber optic bundle which can detect the light emitted by the light sources. In addition, when the transmitter includes an RF transmitter, the receiver includes a directional antenna. In any of the above cases, or similar cases, the relative position between the transmitter and receiver can be determined so that the cutting member 36 of cutting device 32 can be properly located relative to

-9-

vessel 16 such that it re-enters vessel 16 in a region distal of restriction 24.

It should also be noted that, in accordance with another preferred embodiment, vessel 40 is not  
5 formed such that it pierces vessel 16 in the region distal of restriction 24. Rather, vessel 40 is terminated just short of piercing the wall of vessel 16. In that case, blood carrying channels which have intersected the path of lumen 40 through the heart  
10 tissue are allowed to carry the blood to the tissue surrounding vessel 16. Further, the angiogenesis process which is part of the natural healing process can also develop branch vessels, arteriols, and capillaries from the site of lumen 40 to the tissue surrounding  
15 vessel 16, and angiogenesis can be stimulated by the introduction of growth factors.

It should also be noted that, in one preferred embodiment, rather than tunneling around restriction 24, cutting device 31 is advanced through lumen 16 and is  
20 used to form, or drill, orthogonal holes through the vessel wall into any or all of layers 15, 17, 19 and 21. The orthogonal holes provide relief for blood flow through and around restriction 24, and blood carrying channels which have been intersected by the path of the  
25 orthogonal holes through the tissue are allowed to carry the blood to the tissue surrounding vessel 16. Further, the angiogenesis process can also develop branch vessels, arterioles, and capillaries from the site of the orthogonal holes to the tissue surrounding vessel  
30 16. Any suitable cutting device can be used to form the orthogonal holes, such as a mechanical drill-type device, or such as a laser reflected laterally within lumen 16.

-10-

Further, in one embodiment, blood flow through vessel 16 is stopped during the procedure. One system for stopping blood flow includes occluding balloons. Occluding balloons have a fairly low instance of emboli  
5 formation, and therefore have a fairly low instance of neurological problems which result from the formation of emboli.

Thus, it can be seen that, by tunneling through the vessel wall and heart tissue to form a lumen  
10 therethrough which communicates between the proximal and distal ends of lumen 16, around restriction 24, restriction 24 is effectively bypassed. The tunnel can also be formed directly from the aorta to a region in the restricted lumen distal of restriction 24. This is  
15 accomplished without the need for open heart surgery and its associated disadvantages.

Although the present invention has been described with reference to preferred embodiments, workers skilled in the art will recognize that changes  
20 may be made in form and detail without departing from the spirit and scope of the invention.

-11-

WHAT IS CLAIMED IS:

1. A method of bypassing a restriction in a parent vessel lying adjacent heart tissue in a heart, the heart communicating with an aorta, the method comprising:

accessing a first region of the parent vessel  
on a first side of the restriction;

forming a first aperture in the parent vessel  
in the first region;

forming a lumen through the heart tissue, the  
lumen communicating with the parent  
vessel through the first aperture; and

forming a second aperture in a second region  
of the parent vessel on a second side of  
the restriction such that the lumen  
through the heart tissue communicates  
with the parent vessel through the  
second aperture.

2. The method of claim 1 and further comprising:  
delivering a substance to the lumen in the  
heart tissue to increase  
endothelialization in the lumen in the  
heart tissue.

3. The method of claim 2 wherein delivering a  
substance comprises:

delivering a growth factor to the lumen in  
the heart tissue.

4. The method of claim 2 wherein delivering a  
substance comprises:

delivering endothelial cells to the lumen in  
the heart tissue.

5. The method of claim 1 wherein forming the  
first aperture comprises:

-12-

forming the first aperture in a region proximal of the restriction.

6. The method of claim 1 wherein forming a second aperture comprises:

forming the second aperture in a region distal of the restriction, by accessing a wall of the parent vessel from the lumen in the heart tissue.

7. The method of claim 1 wherein accessing a first region of the parent vessel comprises:

advancing a guide catheter through the aorta such that the distal end of the guide catheter lies proximate an ostium of the parent vessel; and

advancing a cutting device through the guide catheter and into the parent vessel proximate the first region in the parent vessel.

8. A method of bypassing a restriction in a parent vessel lying adjacent heart tissue in a heart, the heart communicating with an aorta, the method comprising:

accessing a first region of the parent vessel on a proximal side of the restriction;

forming a first aperture in the parent vessel in the first region;

forming a lumen through the heart tissue, the lumen communicating with the parent vessel through the first aperture, the lumen extending from the first aperture to a region proximate the parent vessel distal of the restriction.

-13-

9. The method of claim 8. and further comprising:  
extending the lumen to form a second aperture  
in a second region of the parent vessel  
distal of the restriction such that the  
lumen through the heart tissue  
communicates with the parent vessel  
through the first and second apertures,  
around the restriction, bypassing the  
restriction.
10. The method of claim 9 wherein accessing a  
first region of the parent vessel comprises:  
advancing a guide catheter through the aorta  
such that the distal end of the guide  
catheter is located proximate an ostium  
of the parent vessel; and  
advancing a cutting device through the guide  
catheter and through the parent vessel  
to the first region of the parent  
vessel.
11. A method of bypassing a restriction in a  
parent vessel lying adjacent heart tissue in a heart,  
the heart communicating with an aorta, the method  
comprising:  
accessing the heart tissue with a cutting  
device;  
forming a lumen through the heart tissue to a  
region proximate the parent vessel  
distal of the restriction; and  
forming an aperture in the parent vessel  
distal of the restriction such that the  
lumen through the heart communicates  
with the aorta and with the parent  
vessel through the aperture in the  
parent vessel.

-14-

12. The method of claim 11 wherein accessing the heart tissue comprises:

accessing a first region of the parent vessel proximal of the restriction; and  
forming a first aperture in the parent vessel in the first region to access the heart tissue.

13. The method of claim 12 wherein accessing a first region of the parent vessel comprises:


advancing a guide catheter through the aorta such that the distal end of the guide catheter is located proximate an ostium of the parent vessel; and

advancing a cutting device through the guide catheter and through the parent vessel to the first region of the parent vessel.

14. The method of claim 11 wherein forming an aperture in the parent vessel distal of the occlusion comprises:

forming the aperture distal of the restriction from within the lumen in the heart tissue.

1/6





2/6






Fig. 1B

3/6

*Fig. 2*

4/6

*Fig. 3*

5/6






Fig. 4

6/6



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/04797

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(6) :A61B 19/00 US CL :128/898 According to International Patent Classification (IPC) or to both national classification and IPC														
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 128/898; 604/49, 96; 606/194; 623/2, 3  Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS, DIALOG														
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>														
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.												
X --- A	US 5,409,019 A (WILK) 25 April 1995, entire document.	8 ----- 1-7, 9-14												
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.														
<table border="0"><tr><td>* Special categories of cited documents:</td><td>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td></tr><tr><td>*A* document defining the general state of the art which is not considered to be of particular relevance</td><td>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td></tr><tr><td>*B* earlier document published on or after the international filing date</td><td>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td></tr><tr><td>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td><td>*Z* document member of the same patent family</td></tr><tr><td>*O* document referring to an oral disclosure, use, exhibition or other means</td><td></td></tr><tr><td>*P* document published prior to the international filing date but later than the priority date claimed</td><td></td></tr></table>			* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	*B* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family	*O* document referring to an oral disclosure, use, exhibition or other means		*P* document published prior to the international filing date but later than the priority date claimed	
* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention													
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone													
*B* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art													
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family													
*O* document referring to an oral disclosure, use, exhibition or other means														
*P* document published prior to the international filing date but later than the priority date claimed														
Date of the actual completion of the international search 13 APRIL 1998		Date of mailing of the international search report 20 MAY 1998												
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer  KELLY O'HARA Telephone No. (703) 308-0780												